

# Computational Intelligence

Winter Term 2009/10

Prof. Dr. Günter Rudolph

Lehrstuhl für Algorithm Engineering (LS 11)

Fakultät für Informatik

TU Dortmund

- Fuzzy sets
  - Axioms of fuzzy complement, t- and s-norms
  - Generators
  - Dual tripels

**Considered so far:**

Standard fuzzy operators

- $A^c(x) = 1 - A(x)$
- $(A \cap B)(x) = \min \{ A(x), B(x) \}$
- $(A \cup B)(x) = \max \{ A(x), B(x) \}$

⇒ Compatible with operators for crisp sets

with membership functions with values in  $\mathbb{B} = \{ 0, 1 \}$

∃ Non-standard operators? ⇒ Yes! Innumerable many!

- Defined via axioms.
- Creation via generators.

### Definition

A function  $c: [0,1] \rightarrow [0,1]$  is a **fuzzy complement** iff

(A1)  $c(0) = 1$  and  $c(1) = 0$ .

(A2)  $\forall a, b \in [0,1]: a \leq b \Rightarrow c(a) \geq c(b)$ .

monotone decreasing

“nice to have”:

(A3)  $c(\cdot)$  is continuous.

(A4)  $\forall a \in [0,1]: c(c(a)) = a$

involution

### Examples:

a) standard fuzzy complement  $c(a) = 1 - a$


ad (A1):  $c(0) = 1 - 0 = 1$  and  $c(1) = 1 - 1 = 0$

ad (A2):  $c'(a) = -1 < 0$  (monotone decreasing)

ad (A3):


ad (A4):  $1 - (1 - a) = a$

$$b) c(a) = \begin{cases} 1 & \text{if } a \leq t \\ 0 & \text{otherwise} \end{cases} \quad \text{for some } t \in (0, 1)$$



ad (A1):  $c(0) = 1$  since  $0 < t$  and  $c(1) = 0$  since  $t < 1$ .

ad (A2): monotone (actually: constant) from 0 to t and t to 1, decreasing at t




ad (A3): **not valid** → discontinuity at t

ad (A4): **not valid** → counter example

$$c(c(1/4)) = c(1) = 0 \neq 1/4 \text{ for } t = 1/2$$

$$c) \ c(a) = \frac{1 + \cos(\pi a)}{2}$$



ad (A1):  $c(0) = 1$  and  $c(1) = 0$

ad (A2):  $c'(a) = -\frac{1}{2} \pi \sin(\pi a) < 0$  since  $\sin(\pi a) > 0$  for  $a \in (0, 1)$



ad (A3): is continuous as a composition of continuous functions

ad (A4): **not valid** → counter example

$$c\left(c\left(\frac{1}{3}\right)\right) = c\left(\frac{3}{4}\right) = \frac{1}{2} \left(1 - \frac{1}{\sqrt{2}}\right) \neq \frac{1}{3}$$

d)  $c(a) = \frac{1-a}{1+\lambda a}$  for  $\lambda > -1$       **Sugeno class**

ad (A1):  $c(0) = 1$  and  $c(1) = 0$

ad (A2):  $c(a) \geq c(b) \Leftrightarrow \frac{1-a}{1+\lambda a} \geq \frac{1-b}{1+\lambda b} \Leftrightarrow$

$(1-a)(1+\lambda b) \geq (1-b)(1+\lambda a) \Leftrightarrow$

$b(\lambda+1) \geq a(\lambda+1) \Leftrightarrow b \geq a$



ad (A3): is continuous as a composition of continuous functions

ad (A4):  $c(c(a)) = c\left(\frac{1-a}{1+\lambda a}\right) = \frac{1-\frac{1-a}{1+\lambda a}}{1+\lambda \frac{1-a}{1+\lambda a}} = \frac{a(\lambda+1)}{\lambda+1} = a$




e)  $c(a) = (1 - a^w)^{1/w}$  for  $w > 0$

**Yager class**


ad (A1):  $c(0) = 1$  and  $c(1) = 0$

ad (A2):  $(1 - a^w)^{1/w} \geq (1 - b^w)^{1/w} \Leftrightarrow 1 - a^w \geq 1 - b^w \Leftrightarrow$   
 $a^w \leq b^w \Leftrightarrow a \leq b$



ad (A3): is continuous as a composition of continuous functions

ad (A4):  $c(c(a)) = c\left(\left(1 - a^w\right)^{\frac{1}{w}}\right) = \left(1 - \left[\left(1 - a^w\right)^{\frac{1}{w}}\right]^w\right)^{\frac{1}{w}}$   
 $= \left(1 - (1 - a^w)\right)^{\frac{1}{w}} = (a^w)^{\frac{1}{w}} = a$






### Theorem


If function  $c: [0,1] \rightarrow [0,1]$  satisfies axioms (A1) and (A2) of fuzzy complement then it has at most one fixed point  $a^*$  with  $c(a^*) = a^*$ .

### Proof:

one fixed point  $\rightarrow$  see example (a)  $\rightarrow$  intersection with bisectrix



no fixed point  $\rightarrow$  see example (b)  $\rightarrow$  no intersection with bisectrix



assume  $\exists n > 1$  fixed points, for example  $a^*$  and  $b^*$  with  $a^* < b^*$

$\Rightarrow c(a^*) = a^*$  and  $c(b^*) = b^*$  (fixed points)

$\Rightarrow c(a^*) < c(b^*)$  with  $a^* < b^*$  impossible if  $c(\cdot)$  is monotone decreasing

$\Rightarrow$  contradiction to axiom (A2) ■

### Theorem

If function  $c: [0, 1] \rightarrow [0, 1]$  satisfies axioms (A1) – (A3) of fuzzy complement then it has exactly one fixed point  $a^*$  with  $c(a^*) = a^*$ .

### Proof:

Intermediate value theorem  $\rightarrow$

If  $c(\cdot)$  continuous (A3) and  $c(0) \geq c(1)$  (A1/A2)

then  $\forall v \in [c(1), c(0)] = [0, 1]: \exists a \in [0, 1]: c(a) = v$ .

$\Rightarrow$  there must be an intersection with bisectrix

$\Rightarrow$  a fixed point exists and by previous theorem there are no other fixed points! ■

### Examples:

$$(a) \quad c(a) = 1 - a \quad \Rightarrow \quad a = 1 - a \quad \Rightarrow \quad a^* = \frac{1}{2}$$

$$(b) \quad c(a) = (1 - a^w)^{1/w} \quad \Rightarrow \quad a = (1 - a^w)^{1/w} \quad \Rightarrow \quad a^* = \left(\frac{1}{2}\right)^{1/w}$$

### Theorem

$c: [0,1] \rightarrow [0,1]$  is involutive fuzzy complement iff

$\exists$  continuous function  $g: [0,1] \rightarrow \mathbb{R}$  with

- $g(0) = 0$
- strictly monotone increasing
- $\forall a \in [0,1]: c(a) = g^{(-1)}(g(1) - g(a))$ . ■

defines an  
**increasing generator**

$g^{(-1)}(x)$  pseudo-inverse

### Examples

a)  $g(x) = x \quad \Rightarrow \quad g^{-1}(x) = x \quad \Rightarrow \quad c(a) = 1 - a \quad \text{(Standard)}$

b)  $g(x) = x^w \quad \Rightarrow \quad g^{-1}(x) = x^{1/w} \quad \Rightarrow \quad c(a) = (1 - a^w)^{1/w} \quad \text{(Yager class, } w > 0)$

c)  $g(x) = \log(x+1) \Rightarrow g^{-1}(x) = e^x - 1 \Rightarrow c(a) = \exp(\log(2) - \log(a+1)) - 1$   
 $= \frac{1 - a}{1 + a} \quad \text{(Sugeno class. } \lambda = 1)$

## Examples

$$d) \quad g(a) = \frac{1}{\lambda} \log_e(1 + \lambda a) \text{ for } \lambda > -1$$

- $g(0) = \log_e(1) = 0$
- strictly monotone increasing since  $g'(a) = \frac{1}{1+\lambda a} > 0$  for  $a \in [0, 1]$
- inverse function on  $[0, 1]$  is  $g^{-1}(a) = \frac{\exp(\lambda a) - 1}{\lambda}$ , thus

$$\begin{aligned} c(a) &= g^{-1} \left( \frac{\log(1 + \lambda)}{\lambda} - \frac{\log(1 + \lambda a)}{\lambda} \right) \\ &= \frac{\exp(\log(1 + \lambda) - \log(1 + \lambda a)) - 1}{\lambda} \\ &= \frac{1}{\lambda} \left( \frac{1 + \lambda}{1 + \lambda a} - 1 \right) = \frac{1 - a}{1 + \lambda a} \quad (\text{Sugeno Complement}) \end{aligned}$$

### Theorem

$c: [0,1] \rightarrow [0,1]$  is involutive fuzzy complement iff

$\exists$  continuous function  $f: [0,1] \rightarrow \mathbb{R}$  with

- $f(1) = 0$
- strictly monotone decreasing
- $\forall a \in [0,1]: c(a) = f^{(-1)}( f(0) - f(a) )$ . ■

defines a  
**decreasing generator**

$f^{(-1)}(x)$  pseudo-inverse

### Examples

$$\text{a) } f(x) = -k - k \cdot x \quad (k > 0) \quad f^{(-1)}(x) = 1 - x/k \quad c(a) = 1 - \frac{-k - (-k - ka)}{k} = 1 - a$$

$$\text{b) } f(x) = 1 - x^w \quad f^{(-1)}(x) = (1 - x)^{1/w} \quad c(a) = f^{-1}(a^w) = (1 - a^w)^{1/w} \quad (\text{Yager})$$

**Definition**

A function  $t: [0, 1] \times [0, 1] \rightarrow [0, 1]$  is a *fuzzy intersection* or *t-norm* iff

$$(A1) \quad t(a, 1) = a$$

$$(A2) \quad b \leq d \Rightarrow t(a, b) \leq t(a, d) \quad (\text{monotonicity})$$

$$(A3) \quad t(a, b) = t(b, a) \quad (\text{commutative})$$

$$(A4) \quad t(a, t(b, d)) = t(t(a, b), d) \quad (\text{associative}) \quad \blacksquare$$

**“nice to have”**

$$(A5) \quad t(a, b) \text{ is continuous} \quad (\text{continuity})$$

$$(A6) \quad t(a, a) < a \quad (\text{subidempotent})$$

$$(A7) \quad a_1 \leq a_2 \text{ and } b_1 \leq b_2 \Rightarrow t(a_1, b_1) \leq t(a_2, b_2) \quad (\text{strict monotonicity})$$

**Note:** the only idempotent t-norm is the standard fuzzy intersection

### Examples:

Name	Function
Standard	$t(a, b) = \min \{ a, b \}$
Algebraic Product	$t(a, b) = a \cdot b$
Bounded Difference	$t(a, b) = \max \{ 0, a + b - 1 \}$
Drastic Product	$t(a, b) = \begin{cases} a & \text{if } b = 1 \\ b & \text{if } a = 1 \\ 0 & \text{otherwise} \end{cases}$

Is algebraic product a t-norm? Check the 4 axioms!

ad (A1):  $t(a, 1) = a \cdot 1 = a$

ad (A3):  $t(a, b) = a \cdot b = b \cdot a = t(b, a)$

ad (A2):  $a \cdot b \leq a \cdot d \Leftrightarrow b \leq d$

ad (A4):  $a \cdot (b \cdot d) = (a \cdot b) \cdot d$

**Theorem**

Function  $t: [0,1] \times [0,1] \rightarrow [0,1]$  is a t-norm  $\Leftrightarrow$

$\exists$  decreasing generator  $f: [0,1] \rightarrow \mathbb{R}$  with  $t(a, b) = f^{(-1)}( f(a) + f(b) )$ . ■

**Example:**

$f(x) = 1/x - 1$  is decreasing generator since

- $f(x)$  is continuous ☑
- $f(1) = 1/1 - 1 = 0$  ☑
- $f'(x) = -1/x^2 < 0$  (monotone decreasing) ☑

inverse function is  $f^{-1}(x) = \frac{1}{x+1}$

$$\Rightarrow t(a, b) = f^{-1} \left( \frac{1}{a} + \frac{1}{b} - 2 \right) = \frac{1}{\frac{1}{a} + \frac{1}{b} - 1} = \frac{ab}{a+b-a-b}$$



**Definition**

A function  $s: [0,1] \times [0,1] \rightarrow [0,1]$  is a *fuzzy union* or *s-norm* or *t-conorm* iff

$$(A1) \quad s(a, 0) = a$$

$$(A2) \quad b \leq d \Rightarrow s(a, b) \leq s(a, d) \quad (\text{monotonicity})$$

$$(A3) \quad s(a, b) = s(b, a) \quad (\text{commutative})$$

$$(A4) \quad s(a, s(b, d)) = s(s(a, b), d) \quad (\text{associative}) \quad \blacksquare$$

**“nice to have”**

$$(A5) \quad s(a, b) \text{ is continuous} \quad (\text{continuity})$$

$$(A6) \quad s(a, a) > a \quad (\text{superidempotent})$$

$$(A7) \quad a_1 \leq a_2 \text{ and } b_1 \leq b_2 \Rightarrow s(a_1, b_1) \leq s(a_2, b_2) \quad (\text{strict monotonicity})$$

**Note:** the only idempotent s-norm is the standard fuzzy union

### Examples:

Name	Function
Standard	$s(a, b) = \max \{ a, b \}$
Algebraic Sum	$s(a, b) = a + b - a \cdot b$
Bounded Sum	$s(a, b) = \min \{ 1, a + b \}$
Drastic Union	$s(a, b) = \begin{cases} a & \text{if } b = 0 \\ b & \text{if } a = 0 \\ 1 & \text{otherwise} \end{cases}$

Is algebraic sum a t-norm? Check the 4 axioms!

ad (A1):  $s(a, 0) = a + 0 - a \cdot 0 = a$

ad (A3):

ad (A2):  $a + b - a \cdot b \leq a + d - a \cdot d \Leftrightarrow b(1 - a) \leq d(1 - a) \Leftrightarrow b \leq d$

ad (A4):

**Theorem**

Function  $s: [0,1] \times [0,1] \rightarrow [0,1]$  is a s-norm  $\Leftrightarrow$

$\exists$  increasing generator  $g: [0,1] \rightarrow \mathbb{R}$  with  $s(a, b) = g^{(-1)}( g(a) + g(b) )$ . ■

**Example:**

$g(x) = -\log(1 - a)$  is decreasing generator since

- $g(x)$  is continuous ☑
- $g(0) = -\log(1 - 0) = 0$  ☑
- $g'(x) = 1/(1-a) > 0$  (monotone increasing) ☑

inverse function is  $g^{-1}(x) = 1 - \exp(-a)$

$$\begin{aligned} \Rightarrow s(a, b) &= g^{-1}(-\log(1 - a) - \log(1 - b)) \\ &= 1 - \exp(\log(1 - a) + \log(1 - b)) \\ &= 1 - (1 - a)(1 - b) = a + b - ab \quad (\text{algebraic sum}) \end{aligned}$$

### Background from classical set theory:

$\cap$  and  $\cup$  operations are dual w.r.t. complement since they obey DeMorgan's laws

#### Definition

A pair of t-norm  $t(\cdot, \cdot)$  and s-norm  $s(\cdot, \cdot)$  is said to be **dual with regard to the fuzzy complement**  $c(\cdot)$  iff

- $c(t(a, b)) = s(c(a), c(b))$
- $c(s(a, b)) = t(c(a), c(b))$

for all  $a, b \in [0, 1]$ . ■

#### Definition

Let  $(c, s, t)$  be a triple of fuzzy complement  $c(\cdot)$ , s- and t-norm.

If  $t$  and  $s$  are dual to  $c$  then the triple  $(c, s, t)$  is called a **dual triple**. ■

### Examples of dual triples

t-norm	s-norm	complement
$\min \{ a, b \}$	$\max \{ a, b \}$	$1 - a$
$a \cdot b$	$a + b - a \cdot b$	$1 - a$
$\max \{ 0, a + b - 1 \}$	$\min \{ 1, a + b \}$	$1 - a$