

Computational Intelligence

Winter Term 2016/17

Prof. Dr. Günter Rudolph

Lehrstuhl für Algorithm Engineering (LS 11)

Fakultät für Informatik

TU Dortmund

- Fuzzy relations
- Fuzzy logic
 - Linguistic variables and terms
 - Inference from fuzzy statements

relations with conventional sets $\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n$:

$$R(\mathcal{X}_1, \mathcal{X}_2, \dots, \mathcal{X}_n) \subseteq \mathcal{X}_1 \times \mathcal{X}_2 \times \dots \times \mathcal{X}_n$$

notice that cartesian product is a **set**!

⇒ all set operations remain valid!

crisp membership function (of x to relation R)

$$R(x_1, x_2, \dots, x_n) = \begin{cases} 1 & \text{if } (x_1, x_2, \dots, x_n) \in R \\ 0 & \text{otherwise} \end{cases}$$

Definition

Fuzzy relation = fuzzy set over crisp cartesian product $\mathcal{X}_1 \times \mathcal{X}_2 \times \dots \times \mathcal{X}_n$ ■

→ each tuple (x_1, \dots, x_n) has a degree of membership to relation


→ degree of membership expresses
strength of relationship between elements of tuple

appropriate representation: n-dimensional membership matrix

example: Let $X = \{ \text{New York}, \text{Paris} \}$ and $Y = \{ \text{Beijing}, \text{New York}, \text{Dortmund} \}$.

relation R = “very far away”

membership matrix



relation R	New York	Paris
Beijing	1.0	0.9
New York	0.0	0.7
Dortmund	0.6	0.3

Definition

Let $R(X, Y)$ be a fuzzy relation with membership matrix R . The **inverse fuzzy relation** to $R(X, Y)$, denoted $R^{-1}(Y, X)$, is a relation on $Y \times X$ with membership matrix $R^{-1} = R'$. ■

Remark: R' is the transpose of membership matrix R .

Evidently: $(R^{-1})^{-1} = R$ since $(R')' = R$

Definition

Let $P(X, Y)$ and $Q(Y, Z)$ be fuzzy relations. The operation \circ on two relations, denoted $P(X, Y) \circ Q(Y, Z)$, is termed **max-min-composition** iff

$$R(x, z) = (P \circ Q)(x, z) = \max_{y \in Y} \min \{ P(x, y), Q(y, z) \}.$$

■

Theorem

- a) max-min composition is associative.
- b) max-min composition is not commutative.
- c) $(P(X,Y) \circ Q(Y,Z))^{-1} = Q^{-1}(Z,Y) \circ P^{-1}(Y,X)$.

membership matrix of max-min composition
determinable via “fuzzy matrix multiplication”: $R = P \circ Q$

fuzzy matrix multiplication $r_{ij} = \max_k \min\{p_{ik}, q_{kj}\}$

crisp matrix multiplication $r_{ij} = \sum_k p_{ik} \cdot q_{kj}$

further methods for realizing compositions of relations:

max-prod composition

$$(P \odot Q)(x, z) = \max_{y \in \mathcal{Y}} \{P(x, y) \cdot Q(y, z)\}$$

generalization: sup-t composition

$$(P \circ Q)(x, z) = \sup_{y \in \mathcal{Y}} \{t(P(x, y), Q(y, z))\}, \text{ where } t(\cdot, \cdot) \text{ is a t-norm}$$

e.g.: $t(a, b) = \min\{a, b\} \Rightarrow$ max-min-composition

$t(a, b) = a \cdot b \Rightarrow$ max-prod-composition

Binary fuzzy relations on $X \times X$: properties

• **reflexive** $\Leftrightarrow \forall x \in X : R(x,x) = 1$

• **irreflexive** $\Leftrightarrow \exists x \in X : R(x,x) < 1$

• **antireflexive** $\Leftrightarrow \forall x \in X : R(x,x) < 1$

• **symmetric** $\Leftrightarrow \forall (x,y) \in X \times X : R(x,y) = R(y,x)$

• **asymmetric** $\Leftrightarrow \exists (x,y) \in X \times X : R(x,y) \neq R(y,x)$

• **antisymmetric** $\Leftrightarrow \forall (x,y) \in X \times X : R(x,y) \neq R(y,x)$

• **transitive** $\Leftrightarrow \forall (x,z) \in X \times X : R(x,z) \geq \max_{y \in Y} \min \{ R(x,y), R(y,z) \}$

• **intransitive** $\Leftrightarrow \exists (x,z) \in X \times X : R(x,z) < \max_{y \in Y} \min \{ R(x,y), R(y,z) \}$

• **antittransitive** $\Leftrightarrow \forall (x,z) \in X \times X : R(x,z) < \max_{y \in Y} \min \{ R(x,y), R(y,z) \}$

actually, here: max-min-transitivity (\rightarrow in general: sup-t-transitivity)

binary fuzzy relation on $X \times X$: example

Let X be the set of all cities in Germany.

Fuzzy relation R is intended to represent the concept of „very close to“.

- $R(x,x) = 1$, since every city is certainly very close to itself.


⇒ **reflexive**

- $R(x,y) = R(y,x)$: if city x is very close to city y , then also vice versa.

⇒ **symmetric**

- $R(\text{Dortmund}, \text{Essen}) = 0.8$
- $R(\text{Essen}, \text{Duisburg}) = 0.7$
- $R(\text{Dortmund}, \text{Duisburg}) = 0.5$
- $R(\text{Dortmund}, \text{Hagen}) = 0.9$

⇒ **intransitive**



crisp:


relation R is equivalence relation \Leftrightarrow R reflexive, symmetric, transitive

fuzzy:

relation R is similarity relation \Leftrightarrow R reflexive, symmetric, (max-min-) transitive

Example:

	a	b	c	d	e	f	g
a	1,0	0,8	0,0	0,4	0,0	0,0	0,0
b	0,8	1,0	0,0	0,4	0,0	0,0	0,0
c	0,0	0,0	1,0	0,0	1,0	0,9	0,5
d	0,4	0,4	0,0	1,0	0,0	0,0	0,0
e	0,0	0,0	1,0	0,0	1,0	0,9	0,5
f	0,0	0,0	0,9	0,0	0,9	1,0	0,5
g	0,0	0,0	0,5	0,0	0,5	0,5	1,0




linguistic variable:

variable that can attain several values of linguistic / verbal nature

e.g.: **color** can attain values **red, green, blue, yellow, ...**

values (red, green, ...) of linguistic variable are called **linguistic terms**

linguistic terms are associated with fuzzy sets



fuzzy proposition

$p: \text{temperature is } \textit{high}$



- LV may be associated with several LT : *high, medium, low, ...*
- *high, medium, low* temperature are fuzzy sets over numerical scale of crisp temperatures
- trueness of fuzzy proposition „temperature is high“ for a given **concrete crisp** temperature value v is interpreted as equal to the degree of membership $\textit{high}(v)$ of the fuzzy set *high*

fuzzy proposition

$p: V \text{ is } F$

linguistic variable (LV) linguistic term (LT)

actually:

$p: V \text{ is } F(v)$

and

$T(p) = F(v)$ for a concrete crisp value v

trueness(p)

establishes connection between *degree of membership* of a fuzzy set and the *degree of trueness* of a fuzzy proposition

fuzzy proposition

p: IF *heating* is *hot*, THEN *energy consumption* is *high*



expresses relation between

- a) temperature of heating and
- b) quantity of energy consumption

p: (*heating*, *energy consumption*) $\in R$

relation

fuzzy proposition

p: IF X is A, THEN Y is B



How can we determine / express degree of trueness $T(p)$?

- For crisp, given values x, y we know $A(x)$ and $B(y)$
- $A(x)$ and $B(y)$ must be processed to single value via relation R
- $R(x, y) = \text{function}(A(x), B(y))$ is fuzzy set over $X \times Y$
- as before: interprete $T(p)$ as degree of membership $R(x,y)$

fuzzy proposition

p: IF X is A, THEN Y is B

A is fuzzy set over X

B is fuzzy set over Y

R is fuzzy set over $X \times Y$

$\forall (x,y) \in X \times Y: R(x, y) = \text{Imp}(A(x), B(y))$

What is $\text{Imp}(\cdot, \cdot)$?

\Rightarrow „appropriate“ fuzzy implication $[0,1] \times [0,1] \rightarrow [0,1]$

assumption: we know an „appropriate“ $\text{Imp}(a,b)$.

How can we determine the degree of trueness $T(p)$?

example:

let $\text{Imp}(a, b) = \min\{ 1, 1 - a + b \}$ and consider fuzzy sets

A:	x_1	x_2	x_3
	0.1	0.8	1.0

B:	y_1	y_2
	0.5	1.0

⇒

R	x_1	x_2	x_3
y_1	1.0	0.7	0.5
y_2	1.0	1.0	1.0

z.B.

$$R(x_2, y_1) = \text{Imp}(A(x_2), B(y_1)) = \text{Imp}(0.8, 0.5) = \min\{1.0, 0.7\} = 0.7$$

and $T(p)$ for (x_2, y_1) is $R(x_2, y_1) = 0.7$

■

toward inference from fuzzy statements:



- let $\forall x, y: y = f(x)$.

IF $X = x_0$ THEN $Y = f(x_0)$

- IF $X \in A$ THEN $Y \in B = \{ y \in \mathcal{Y} : y = f(x), x \in A \}$



crisp case:
functional
relationship





toward inference from fuzzy statements:

- let relationship between x and y be a relation R on $\mathcal{X} \times \mathcal{Y}$

$$\text{IF } X = x_0 \text{ THEN } Y \in B = \{ y \in \mathcal{Y} : (x_0, y) \in R \}$$

- IF $X \in A$ THEN $Y \in B = \{ y \in \mathcal{Y} : (x, y) \in R, x \in A \}$

crisp case:
relational
relationship



toward inference from fuzzy statements:


IF $X \in A$ THEN $Y \in B = \{ y \in \mathcal{Y} : (x, y) \in R, x \in A \}$

also expressible via characteristic functions of sets A, B, R :

$$B(y) = 1 \text{ iff } \exists x: A(x) = 1 \text{ and } R(x, y) = 1$$

$$\Leftrightarrow \exists x: \min\{ A(x), R(x, y) \} = 1$$

$$\Leftrightarrow \max_{x \in \mathcal{X}} \min\{ A(x), R(x, y) \} = 1$$



$$\forall y \in \mathcal{Y}: B(y) = \max_{x \in \mathcal{X}} \min\{ A(x), R(x, y) \}$$

inference from fuzzy statements

Now: A' , B' fuzzy sets over \mathcal{X} resp. \mathcal{Y}

Assume: $R(x,y)$ and $A'(x)$ are given.

Idea: Generalize characteristic function of $B(y)$ to membership function $B'(y)$

$$\forall y \in \mathcal{Y}: B(y) = \max_{x \in \mathcal{X}} \min \{ A(x), R(x, y) \}$$

characteristic functions

$$\forall y \in \mathcal{Y}: B'(y) = \sup_{x \in \mathcal{X}} \min \{ A'(x), R(x, y) \}$$

membership functions

composition rule of inference (in matrix form): $B^T = A \circ R$

inference from fuzzy statements

- conventional:
modus ponens

$$\begin{array}{c} a \Rightarrow b \\ a \\ \hline b \end{array}$$

- fuzzy:
generalized modus ponens (GMP)

$$\begin{array}{c} \text{IF } X \text{ is A, THEN } Y \text{ is B} \\ X \text{ is A'} \\ \hline Y \text{ is B'} \end{array}$$

e.g.: IF *heating* is hot, THEN *energy consumption* is high
heating is warm
—————
energy consumption is normal

example: GMP

consider

A:	<table border="1"><tr><td>x_1</td><td>x_2</td><td>x_3</td></tr><tr><td>0.5</td><td>1.0</td><td>0.6</td></tr></table>	x_1	x_2	x_3	0.5	1.0	0.6
x_1	x_2	x_3					
0.5	1.0	0.6					

B:	<table border="1"><tr><td>y_1</td><td>y_2</td></tr><tr><td>1.0</td><td>0.4</td></tr></table>	y_1	y_2	1.0	0.4
y_1	y_2				
1.0	0.4				

with the rule: IF X is A THEN Y is B

given fact

A':	<table border="1"><tr><td>x_1</td><td>x_2</td><td>x_3</td></tr><tr><td>0.6</td><td>0.9</td><td>0.7</td></tr></table>	x_1	x_2	x_3	0.6	0.9	0.7
x_1	x_2	x_3					
0.6	0.9	0.7					

 \Rightarrow

R	x_1	x_2	x_3
y_1	1.0	1.0	1.0
y_2	0.9	0.4	0.8

with $\text{Imp}(a,b) = \min\{1, 1-a+b\}$ thus: $A' \circ R = B'$

with max-min-composition

$$\begin{pmatrix} 0.6 & 0.9 & 0.7 \end{pmatrix} \circ \begin{pmatrix} 1.0 & 0.9 \\ 1.0 & 0.4 \\ 1.0 & 0.8 \end{pmatrix} = \begin{pmatrix} 0.9 & 0.7 \end{pmatrix}$$

inference from fuzzy statements

- conventional:
modus tollens

$$\frac{a \Rightarrow b}{\frac{\overline{b}}{\overline{a}}}$$

- fuzzy:
generalized modus tollens (GMT)

$$\frac{\text{IF } X \text{ is } A, \text{ THEN } Y \text{ is } B \\ Y \text{ is } B'}{X \text{ is } A'}$$

e.g.: IF *heating* is hot, THEN *energy consumption* is high
energy consumption is normal
heating is warm

example: GMT

consider

A:

x_1	x_2	x_3
0.5	1.0	0.6

B:

y_1	y_2
1.0	0.4

with the rule: IF X is A THEN Y is B

given fact

B':

y_1	y_2
0.9	0.7

⇒

R	x_1	x_2	x_3
y_1	1.0	1.0	1.0
y_2	0.9	0.4	0.8

with $\text{Imp}(a,b) = \min\{1, 1-a+b\}$

thus: $B' \circ R^{-1} = A'$ $(0.9 \ 0.7) \circ \begin{pmatrix} 1.0 & 1.0 & 1.0 \\ 0.9 & 0.4 & 0.8 \end{pmatrix} = (0.9 \ 0.9 \ 0.9)$

with max-min-composition

inference from fuzzy statements

- conventional:
hypothetic syllogism

$$\begin{array}{c} a \Rightarrow b \\ b \Rightarrow c \\ \hline a \Rightarrow c \end{array}$$

- fuzzy:
generalized HS

$$\begin{array}{c} \text{IF } X \text{ is A, THEN } Y \text{ is B} \\ \text{IF } Y \text{ is B, THEN } Z \text{ is C} \\ \hline \text{IF } X \text{ is A, THEN } Z \text{ is C} \end{array}$$

e.g.: IF *heating* is hot, THEN *energy consumption* is high
 IF *energy consumption* is high, THEN *living* is expensive

 IF *heating* is hot, THEN *living* is expensive

example: GHS

let fuzzy sets $A(x)$, $B(y)$, $C(z)$ be given

⇒ determine the three relations

$$R_1(x,y) = \text{Imp}(A(x),B(y))$$

$$R_2(y,z) = \text{Imp}(B(y),C(z))$$

$$R_3(x,z) = \text{Imp}(A(x),C(z))$$

and express them as matrices R_1 , R_2 , R_3

We say:

GHS is valid if $R_1 \circ R_2 = R_3$

So, ... what makes sense for $\text{Imp}(\cdot, \cdot)$?

$\text{Imp}(a, b)$ ought to express fuzzy version of implication $(a \Rightarrow b)$

conventional: $a \Rightarrow b$ identical to $\bar{a} \vee b$

But how can we calculate with fuzzy “boolean” expressions?

request: must be compatible to crisp version (and more) for $a, b \in \{0, 1\}$

a	b	$a \wedge b$	$t(a, b)$
0	0	0	0
0	1	0	0
1	0	0	0
1	1	1	1

a	b	$a \vee b$	$s(a, b)$
0	0	0	0
0	1	1	1
1	0	1	1
1	1	1	1

a	\bar{a}	$c(a)$
0	1	1
1	0	0

So, ... what makes sense for $\text{Imp}(\cdot, \cdot)$?

1st approach: S implications

conventional: $a \Rightarrow b$ identical to $\bar{a} \vee b$

fuzzy: $\text{Imp}(a, b) = s(c(a), b)$

2nd approach: R implications

conventional: $a \Rightarrow b$ identical to $\max\{x \in \{0, 1\} : a \wedge x \leq b\}$

fuzzy: $\text{Imp}(a, b) = \max\{x \in [0, 1] : t(a, x) \leq b\}$

3rd approach: QL implications

conventional: $a \Rightarrow b$ identical to $\bar{a} \vee b \equiv \bar{a} \vee (a \wedge b)$ law of absorption

fuzzy: $\text{Imp}(a, b) = s(c(a), t(a, b))$ (dual tripel ?)

example: S implication

$$\text{Imp}(a, b) = s(c_s(a), b) \quad (c_s : \text{std. complement})$$

1. Kleene-Dienes implication

$$s(a, b) = \max\{ a, b \} \quad (\text{standard})$$

$$\text{Imp}(a, b) = \max\{ 1-a, b \}$$

2. Reichenbach implication

$$s(a, b) = a + b - ab \quad (\text{algebraic sum})$$

$$\text{Imp}(a, b) = 1 - a + ab$$

3. Łukasiewicz implication

$$s(a, b) = \min\{ 1, a + b \} \quad (\text{bounded sum})$$

$$\text{Imp}(a, b) = \min\{ 1, 1 - a + b \}$$

example: R implicationen

$$\text{Imp}(a, b) = \max\{ x \in [0, 1] : t(a, x) \leq b \}$$

1. Gödel implication

$$t(a, b) = \min\{ a, b \}$$

(std.)

$$\text{Imp}(a, b) = \begin{cases} 1 & , \text{ if } a \leq b \\ b & , \text{ else} \end{cases}$$

2. Goguen implication

$$t(a, b) = ab$$

(algeb. product)

$$\text{Imp}(a, b) = \begin{cases} 1 & , \text{ if } a \leq b \\ \frac{b}{a} & , \text{ else} \end{cases}$$

3. Łukasiewicz implication

$$t(a, b) = \max\{ 0, a + b - 1 \} \quad (\text{bounded diff.})$$

$$\text{Imp}(a, b) = \min\{ 1, 1 - a + b \}$$

example: QL implication

$$\text{Imp}(a, b) = s(c(a), t(a, b))$$

1. Zadeh implication

$$\begin{aligned} t(a, b) &= \min \{ a, b \} && \text{(std.)} \\ s(a, b) &= \max \{ a, b \} && \text{(std.)} \end{aligned}$$

$$\text{Imp}(a, b) = \max \{ 1 - a, \min \{ a, b \} \}$$

2. „NN“ implication ☺ (Klir/Yuan 1994)

$$\begin{aligned} t(a, b) &= ab && \text{(algebr. prd.)} \\ s(a, b) &= a + b - ab && \text{(algebr. sum)} \end{aligned}$$

$$\text{Imp}(a, b) = 1 - a + a^2b$$

3. Kleene-Dienes implication

$$\begin{aligned} t(a, b) &= \max \{ 0, a + b - 1 \} && \text{(bounded diff.)} \\ s(a, b) &= \min \{ 1, a + b \} && \text{(bounded sum)} \end{aligned}$$

axioms for fuzzy implications

- | | |
|--|--------------------------|
| 1. $a \leq b$ implies $\text{Imp}(a, x) \geq \text{Imp}(b, x)$ | monotone in 1st argument |
| 2. $a \leq b$ implies $\text{Imp}(x, a) \leq \text{Imp}(x, b)$ | monotone in 2nd argument |
| 3. $\text{Imp}(0, a) = 1$ | dominance of falseness |
| 4. $\text{Imp}(1, b) = b$ | neutrality of trueness |
| 5. $\text{Imp}(a, a) = 1$ | identity |
| 6. $\text{Imp}(a, \text{Imp}(b, x)) = \text{Imp}(b, \text{Imp}(a, x))$ | exchange property |
| 7. $\text{Imp}(a, b) = 1$ iff $a \leq b$ | boundary condition |
| 8. $\text{Imp}(a, b) = \text{Imp}(c(b), c(a))$ | contraposition |
| 9. $\text{Imp}(\cdot, \cdot)$ is continuous | continuity |

characterization of fuzzy implication

Theorem:

$\text{Imp}: [0,1] \times [0,1] \rightarrow [0,1]$ satisfies axioms 1-9 for fuzzy implications
for a certain fuzzy complement $c(\cdot)$ \Leftrightarrow

\exists strictly monotone increasing, continuous function $f: [0,1] \rightarrow [0, \infty)$ with

- $f(0) = 0$
- $\forall a, b \in [0,1]: \text{Imp}(a, b) = f^{-1}(\min\{f(1) - f(a) + f(b), f(1)\})$
- $\forall a \in [0,1]: c(a) = f^{-1}(f(1) - f(a))$

Proof: Smets & Magrez (1987), p. 337f.

■

examples: (in tutorial)

choosing an „appropriate“ fuzzy implication ...

apt quotation: (Klir & Yuan 1995, p. 312)

„To select an appropriate fuzzy implication for approximate reasoning under each particular situation is a difficult problem.“

guideline:

GMP, GMT, GHS should be compatible with MP, MT, HS

for fuzzy implication in calculations with relations:

$$B(y) = \sup \{ t(A(x), \text{Imp}(A(x), B(y))) : x \in X \}$$

example:

Gödel implication for t-norm = bounded difference