

Computational Intelligence

Winter Term 2022/23

Prof. Dr. Günter Rudolph

Lehrstuhl für Algorithm Engineering (LS 11)

Fakultät für Informatik

TU Dortmund

Plan for Today

Lecture 02

- Fuzzy sets
 - Axioms of fuzzy complement, t- and s-norms
 - Generators
 - Dual tripels

Fuzzy Sets

Lecture 02

Considered so far:

Standard fuzzy operators

- $A^c(x) = 1 - A(x)$
- $(A \cap B)(x) = \min \{ A(x), B(x) \}$
- $(A \cup B)(x) = \max \{ A(x), B(x) \}$

⇒ Compatible with operators for crisp sets

with membership functions with values in $\mathbb{B} = \{ 0, 1 \}$

∃ Non-standard operators? ⇒ Yes! Innumerable many!

- Defined via axioms.
- Creation via generators.

Fuzzy Complement: Axioms

Lecture 02

Definition

A function $c: [0,1] \rightarrow [0,1]$ is a **fuzzy complement** iff

$$(A1) \quad c(0) = 1 \text{ and } c(1) = 0.$$

$$(A2) \quad \forall a, b \in [0,1]: a \leq b \Rightarrow c(a) \geq c(b).$$

monotone decreasing

“nice to have”:

$$(A3) \quad c(\cdot) \text{ is continuous.}$$

$$(A4) \quad \forall a \in [0,1]: c(c(a)) = a$$

involutive

Examples:

a) standard fuzzy complement $c(a) = 1 - a$

$$\text{ad (A1): } c(0) = 1 - 0 = 1 \text{ and } c(1) = 1 - 1 = 0$$

$$\text{ad (A2): } c'(a) = -1 < 0 \text{ (monotone decreasing)}$$


ad (A3):

ad (A4): $1 - (1 - a) = a$

Fuzzy Complement: Examples

Lecture 02

b) $c(a) = \begin{cases} 1 & \text{if } a \leq t \\ 0 & \text{otherwise} \end{cases}$ for some $t \in (0, 1)$



ad (A1): $c(0) = 1$ since $0 < t$ and $c(1) = 0$ since $t < 1$.

ad (A2): monotone (actually: constant) from 0 to t and t to 1, decreasing at t

ad (A3): **not valid** → discontinuity at t


ad (A4): **not valid** → counter example

$$c(c(\frac{1}{4})) = c(1) = 0 \neq \frac{1}{4} \text{ for } t = \frac{1}{2}$$

Fuzzy Complement: Examples

Lecture 02

c) $c(a) = \frac{1 + \cos(\pi a)}{2}$



ad (A1): $c(0) = 1$ and $c(1) = 0$

ad (A2): $c'(a) = -\frac{1}{2} \pi \sin(\pi a) < 0$ since $\sin(\pi a) > 0$ for $a \in (0, 1)$

ad (A3): is continuous as a composition of continuous functions;
alternative argument: derivative exists, see $c'(a)$ in (A2)


ad (A4): **not valid** → counter example

$$c(c(\frac{1}{3})) = c(\frac{3}{4}) = \frac{1}{2} \left(1 - \frac{1}{\sqrt{2}}\right) \neq \frac{1}{3}$$

Fuzzy Complement: Examples

Lecture 02

d) $c(a) = \frac{1-a}{1+\lambda a}$ for $\lambda > -1$ **Sugeno class**



ad (A1): $c(0) = 1$ and $c(1) = 0$

$$\begin{aligned} \text{ad (A2): } c(a) \geq c(b) &\Leftrightarrow \frac{1-a}{1+\lambda a} \geq \frac{1-b}{1+\lambda b} \Leftrightarrow \\ &(1-a)(1+\lambda b) \geq (1-b)(1+\lambda a) \Leftrightarrow \\ &b(\lambda+1) \geq a(\lambda+1) \Leftrightarrow b \geq a \end{aligned}$$

ad (A3): is continuous as a composition of continuous functions


$$\text{ad (A4): } c(c(a)) = c\left(\frac{1-a}{1+\lambda a}\right) = \frac{1-\frac{1-a}{1+\lambda a}}{1+\lambda \frac{1-a}{1+\lambda a}} = \frac{a(\lambda+1)}{\lambda+1} = a$$

Fuzzy Complement: Examples

Lecture 02

e) $c(a) = (1 - a^w)^{1/w}$ for $w > 0$

Yager class



ad (A1): $c(0) = 1$ and $c(1) = 0$

$$\begin{aligned} \text{ad (A2): } (1 - a^w)^{1/w} \geq (1 - b^w)^{1/w} &\Leftrightarrow 1 - a^w \geq 1 - b^w \Leftrightarrow \\ &a^w \leq b^w \Leftrightarrow a \leq b \end{aligned}$$

ad (A3): is continuous as a composition of continuous functions


$$\begin{aligned} \text{ad (A4): } c(c(a)) &= c\left((1 - a^w)^{\frac{1}{w}}\right) = \left(1 - \left[(1 - a^w)^{\frac{1}{w}}\right]^w\right)^{\frac{1}{w}} \\ &= (1 - (1 - a^w))^{\frac{1}{w}} = (a^w)^{\frac{1}{w}} = a \end{aligned}$$

Theorem


If function $c:[0,1] \rightarrow [0,1]$ satisfies axioms (A1) and (A2) of fuzzy complement then it has at most one fixed point a^* with $c(a^*) = a^*$.

Proof:

one fixed point \rightarrow see example (a) \rightarrow intersection with bisectrix



no fixed point \rightarrow see example (b) \rightarrow no intersection with bisectrix



assume $\exists n > 1$ fixed points, for example a^* and b^* with $a^* < b^*$

$\Rightarrow c(a^*) = a^*$ and $c(b^*) = b^*$ (fixed points)

$\Rightarrow c(a^*) < c(b^*)$ with $a^* < b^*$ impossible if $c(\cdot)$ is monotone decreasing

\Rightarrow contradiction to axiom (A2) ■

Theorem

If function $c:[0,1] \rightarrow [0,1]$ satisfies axioms (A1) – (A3) of fuzzy complement then it has exactly one fixed point a^* with $c(a^*) = a^*$.

Proof:

Intermediate value theorem \rightarrow

If $c(\cdot)$ continuous (A3) and $c(0) \geq c(1)$ (A1/A2)

then $\forall v \in [c(1), c(0)] = [0,1]: \exists a \in [0,1]: c(a) = v$.

\Rightarrow there must be an intersection with bisectrix

\Rightarrow a fixed point exists and by previous theorem there are no other fixed points! ■

Examples:

$$(a) c(a) = 1 - a \quad \Rightarrow a = 1 - a \quad \Rightarrow a^* = \frac{1}{2}$$

$$(b) c(a) = (1 - a^w)^{1/w} \quad \Rightarrow a = (1 - a^w)^{1/w} \quad \Rightarrow a^* = (\frac{1}{2})^{1/w}$$

Theorem

$c: [0,1] \rightarrow [0,1]$ is involutive fuzzy complement iff

\exists continuous function $g: [0,1] \rightarrow \mathbb{R}$ with

- $g(0) = 0$
- strictly monotone increasing
- $\forall a \in [0,1]: c(a) = g^{-1}(g(1) - g(a))$. ■

defines an
increasing generator
 $g^{-1}(x)$ pseudo-inverse
 $= g^{-1}(\min\{g(1), x\})$

Examples

$$a) g(x) = x \quad \Rightarrow g^{-1}(x) = x \quad \Rightarrow c(a) = 1 - a \quad (\text{Standard})$$

$$b) g(x) = x^w \quad \Rightarrow g^{-1}(x) = x^{1/w} \quad \Rightarrow c(a) = (1 - a^w)^{1/w} \quad (\text{Yager class, } w > 0)$$

$$c) g(x) = \log(x+1) \Rightarrow g^{-1}(x) = e^x - 1 \Rightarrow c(a) = \exp(\log(2) - \log(a+1)) - 1$$

$\underbrace{\quad}_{?} = \frac{1-a}{1+a}$ (Sugeno class. $\lambda = 1$)

\rightarrow make sure that pseudoinverse is equal to inverse, here!

$$g(x) = \log(x+1) \quad \rightarrow \quad g^{-1}(x) = e^x - 1 \quad \checkmark \quad (\text{inverse})$$

$$g^{(-1)}(x) = g^{-1}(\min\{g(1), x\}) \quad ? \quad (\text{pseudoinverse})$$

$$c(a) = g^{(-1)}(\underbrace{g(1) - g(a)}_{= x})$$

$$\min\{g(1), g(1) - a\} = g(1) - g(a) \leq g(1) \text{ since } 0 \leq g(a) \leq \log 2 \text{ for } a \in [0, 1]$$

therefore,

$$c(a) = g^{(-1)}(g(1) - g(a)) = g^{-1}(g(1) - g(a)) \quad \checkmark$$

Examples

d) $g(a) = \frac{1}{\lambda} \log_e(1 + \lambda a)$ for $\lambda > -1$

- $g(0) = \log_e(1) = 0$
- strictly monotone increasing since $g'(a) = \frac{1}{1+\lambda a} > 0$ for $a \in [0, 1]$
- inverse function on $[0, 1]$ is $g^{-1}(a) = \frac{\exp(\lambda a) - 1}{\lambda}$, thus

$$\begin{aligned} c(a) &= g^{-1}\left(\frac{\log(1 + \lambda)}{\lambda} - \frac{\log(1 + \lambda a)}{\lambda}\right) \\ &= \frac{\exp(\log(1 + \lambda)) - \log(1 + \lambda a) - 1}{\lambda} \\ &= \frac{1}{\lambda} \left(\frac{1 + \lambda}{1 + \lambda a} - 1 \right) = \frac{1 - a}{1 + \lambda a} \quad (\text{Sugeno Complement}) \end{aligned}$$

Definition

A function $t:[0,1] \times [0,1] \rightarrow [0,1]$ is a **fuzzy intersection** or **t-norm** iff $\forall a,b,d \in [0,1]$

- | | |
|--------------------------------------------------|----------------------|
| (A1) $t(a, 1) = a$ | (boundary condition) |
| (A2) $b \leq d \Rightarrow t(a, b) \leq t(a, d)$ | (monotonicity) |
| (A3) $t(a, b) = t(b, a)$ | (commutative) |
| (A4) $t(a, t(b, d)) = t(t(a, b), d)$ | (associative) ■ |

“nice to have”

- | | |
|---------------------------------------------------------------------------|-----------------------|
| (A5) $t(a, b)$ is continuous | (continuity) |
| (A6) $t(a, a) < a$ for $0 < a < 1$ | (subidempotent) |
| (A7) $a_1 < a_2$ and $b_1 \leq b_2 \Rightarrow t(a_1, b_1) < t(a_2, b_2)$ | (strict monotonicity) |

Note: the only idempotent t-norm is the standard fuzzy intersection

Theorem

$c: [0,1] \rightarrow [0,1]$ is involutive fuzzy complement iff

\exists continuous function $f: [0,1] \rightarrow \mathbb{R}$ with

- $f(1) = 0$
- strictly monotone decreasing
- $\forall a \in [0,1]: c(a) = f^{-1}(f(0) - f(a))$. ■

defines a
decreasing generator
 $f^{-1}(x)$ pseudo-inverse
 $= f^{-1}(\min\{f(0), x\})$

Examples

a) $f(x) = k - k \cdot x$ ($k \geq 1$) $f^{-1}(x) = 1 - x/k$ $c(a) = 1 - \frac{k-(k-ka)}{k} = 1 - a$

b) $f(x) = 1 - x^w$ $f^{-1}(x) = (1-x)^{1/w}$ $c(a) = f^{-1}(a^w) = (1-a^w)^{1/w}$ (Yager)

Definition

A function $t:[0,1] \times [0,1] \rightarrow [0,1]$ is a **fuzzy intersection** or **t-norm** iff $\forall a,b,d \in [0,1]$

- | | |
|--------------------------------------------------|----------------------|
| (A1) $t(a, 1) = a$ | (boundary condition) |
| (A2) $b \leq d \Rightarrow t(a, b) \leq t(a, d)$ | (monotonicity) |
| (A3) $t(a, b) = t(b, a)$ | (commutative) |
| (A4) $t(a, t(b, d)) = t(t(a, b), d)$ | (associative) ■ |

“nice to have”

- | | |
|---------------------------------------------------------------------------|-----------------------|
| (A5) $t(a, b)$ is continuous | (continuity) |
| (A6) $t(a, a) < a$ for $0 < a < 1$ | (subidempotent) |
| (A7) $a_1 < a_2$ and $b_1 \leq b_2 \Rightarrow t(a_1, b_1) < t(a_2, b_2)$ | (strict monotonicity) |

Note: the only idempotent t-norm is the standard fuzzy intersection

Theorem:

The only idempotent t-norm is the standard fuzzy intersection.

Proof:

Assume there exists a t-norm with $t(a,a) = a$ for all $a \in [0,1]$.

- If $0 \leq a \leq b \leq 1$ then

$$a = t(a,a) \stackrel{\text{by assumption}}{\leq} t(a,b) \stackrel{\text{by monotonicity}}{\leq} t(a,1) = a \stackrel{\text{by boundary condition}}{\leq}$$

and hence $t(a,b) = a$.

- If $0 \leq b \leq a \leq 1$ then

$$b = t(b,b) \stackrel{\text{by assumption}}{\leq} t(b,a) \stackrel{\text{by monotonicity}}{\leq} t(b,1) = b \stackrel{\text{by boundary condition}}{\leq}$$

and hence $t(a,b) = t(b,a) = b$.


↑
by commutativity

$t(a,b) = \min(a,b)$
is the only
possible solution!

q.e.d.

Examples:

Name	Function
(a) Standard	$t(a, b) = \min \{ a, b \}$
(b) Algebraic Product	$t(a, b) = a \cdot b$
(c) Bounded Difference	$t(a, b) = \max \{ 0, a + b - 1 \}$
(d) Drastic Product	$t(a, b) = \begin{cases} a & \text{if } b = 1 \\ b & \text{if } a = 1 \\ 0 & \text{otherwise} \end{cases}$



Is algebraic product a t-norm? Check the 4 axioms!

- | | | | |
|--------------------------------------------------------------|-------------------------------------|------------------------------------------------------|-------------------------------------|
| ad (A1): $t(a, 1) = a \cdot 1 = a$ | <input checked="" type="checkbox"/> | ad (A3): $t(a, b) = a \cdot b = b \cdot a = t(b, a)$ | <input checked="" type="checkbox"/> |
| ad (A2): $a \cdot b \leq a \cdot d \Leftrightarrow b \leq d$ | <input checked="" type="checkbox"/> | ad (A4): $a \cdot (b \cdot d) = (a \cdot b) \cdot d$ | <input checked="" type="checkbox"/> |

Definition

A function $s:[0,1] \times [0,1] \rightarrow [0,1]$ is a **fuzzy union** or **s-norm** iff $\forall a,b,d \in [0,1]$

- | | |
|--------------------------------------------------|----------------------|
| (A1) $s(a, 0) = a$ | (boundary condition) |
| (A2) $b \leq d \Rightarrow s(a, b) \leq s(a, d)$ | (monotonicity) |
| (A3) $s(a, b) = s(b, a)$ | (commutative) |
| (A4) $s(a, s(b, d)) = s(s(a, b), d)$ | (associative) |

“nice to have”

- | | |
|---------------------------------------------------------------------------|-----------------------|
| (A5) $s(a, b)$ is continuous | (continuity) |
| (A6) $s(a, a) > a$ for $0 < a < 1$ | (superidempotent) |
| (A7) $a_1 < a_2$ and $b_1 \leq b_2 \Rightarrow s(a_1, b_1) < s(a_2, b_2)$ | (strict monotonicity) |

Note: the only idempotent s-norm is the standard fuzzy union

Theorem

Function $t: [0,1] \times [0,1] \rightarrow [0,1]$ is a t-norm ,

\exists decreasing generator $f:[0,1] \rightarrow \mathbb{R}$ with $t(a, b) = f^{-1}(\min\{f(0), f(a) + f(b)\})$. ■

Example:

$f(x) = 1/x - 1$ is decreasing generator since


- $f(x)$ is continuous
- $f(1) = 1/1 - 1 = 0$
- $f'(x) = -1/x^2 < 0$ (monotone decreasing)

inverse function is $f^{-1}(x) = \frac{1}{x+1}$; $f(0) = \infty \Rightarrow \min\{f(0), f(a) + f(b)\} = f(a) + f(b)$

$$\Rightarrow t(a, b) = f^{-1}\left(\frac{1}{a} + \frac{1}{b} - 2\right) = \frac{1}{\frac{1}{a} + \frac{1}{b} - 1} = \frac{ab}{a+b-ab}$$

Fuzzy Union: s-norm**Examples:**

Name	Function
Standard	$s(a, b) = \max \{ a, b \}$
Algebraic Sum	$s(a, b) = a + b - a \cdot b$
Bounded Sum	$s(a, b) = \min \{ 1, a + b \}$
Drastic Union	$s(a, b) = \begin{cases} a & \text{if } b = 0 \\ b & \text{if } a = 0 \\ 1 & \text{otherwise} \end{cases}$



Is algebraic sum an s-norm? Check the 4 axioms!

- | | | |
|-----------------------------------------------------------------------------------------------------------------|-------------------------------------|----------------------------------------------|
| ad (A1): $s(a, 0) = a + 0 - a \cdot 0 = a$ | <input checked="" type="checkbox"/> | ad (A3): <input checked="" type="checkbox"/> |
| ad (A2): $a + b - a \cdot b \leq a + d - a \cdot d \Leftrightarrow b(1-a) \leq d(1-a) \Leftrightarrow b \leq d$ | <input checked="" type="checkbox"/> | ad (A4): <input checked="" type="checkbox"/> |

Theorem

Function $s: [0,1] \times [0,1] \rightarrow [0,1]$ is a s-norm \Leftrightarrow

\exists increasing generator $g: [0,1] \rightarrow \mathbb{R}$ with $s(a, b) = g^{-1}(\min\{g(1), g(a) + g(b)\})$. ■


Example:

$g(x) = -\log(1-x)$ is increasing generator since


- $g(x)$ is continuous
- $g(0) = -\log(1-0) = 0$
- $g'(x) = 1/(1-x) > 0$ (monotone increasing)

inverse function is $g^{-1}(x) = 1 - \exp(-x)$; $g(1) = \infty \Rightarrow \min\{g(1), g(a) + g(b)\} = g(a) + g(b)$

$$\begin{aligned} \Rightarrow s(a, b) &= g^{-1}(-\log(1-a) - \log(1-b)) \\ &= 1 - \exp(\log(1-a) + \log(1-b)) \\ &= 1 - (1-a)(1-b) = a+b-ab \quad (\text{algebraic sum}) \end{aligned}$$



$c(t(a, b))$



$s(c(a), c(b))$


Dual Triple:

- bounded difference
- bounded sum
- standard complement

\Rightarrow left image = right image



$c(t(a, b))$



Non-Dual Triple:

- algebraic product
- bounded sum
- standard complement

\Rightarrow left image \neq right image

Background from classical set theory:

\cap and \cup operations are dual w.r.t. complement since they obey DeMorgan's laws

Definition

A pair of t-norm $t(\cdot, \cdot)$ and s-norm $s(\cdot, \cdot)$ is said to be **dual with regard to the fuzzy complement** $c(\cdot)$ iff

- $c(t(a, b)) = s(c(a), c(b))$
- $c(s(a, b)) = t(c(a), c(b))$

for all $a, b \in [0,1]$. ■

Definition

Let (c, s, t) be a triple of fuzzy complement $c(\cdot)$, s- and t-norm.

If t and s are dual to c then the triple (c, s, t) is called a **dual triple**. ■

Examples of dual tripels

t-norm	s-norm	complement
$\min\{a, b\}$	$\max\{a, b\}$	$1-a$
$a \cdot b$	$a+b-a \cdot b$	$1-a$
$\max\{0, a+b-1\}$	$\min\{1, a+b\}$	$1-a$

Why are dual triples so important?

\Rightarrow allow equivalence transformations of fuzzy set expressions

\Rightarrow required to transform into some equivalent normal form (standardized input)

\Rightarrow e.g. two stages: intersection of unions

$$\bigcap_{i=1}^n (A_i \cup B_i)$$

or union of intersections

$$\bigcup_{i=1}^n (A_i \cap B_i)$$

Example:

$$A \cup (B \cap (C \cap D)^c) =$$

\leftarrow not in normal form

$$A \cup (B \cap (C^c \cup D^c)) =$$

\leftarrow equivalent if DeMorgan's law valid (dual triples!)

$$A \cup (B \cap C^c) \cup (B \cap D^c)$$

\leftarrow equivalent (distributive lattice!)